

Dynamic Susceptibility and Phonon Anomalies in the Bilayer t - J Model

B. Normand, H. Kohno and H. Fukuyama

Department of Physics, University of Tokyo,
7-3-1 Hongo, Bunkyo-ku, Tokyo 113, Japan.

We consider a bilayer version of the extended t - J model, with a view to computing the form of certain experimentally observable properties. Using the slave-boson decomposition, we show at the mean-field level that in the bilayer system the existence of in-plane d -wave singlet pairing excludes any interplane singlet order for reasonable values of the interplane superexchange parameter. Restricting the analysis to the regime of no interplane singlet pairing, we deduce parameter sets reproducing the Fermi surfaces of YBCO- and BSCCO-like bilayer systems. From these we calculate the form of the dynamic susceptibility $\chi(\mathbf{q}, \omega)$ in both systems, and of the anomalies in frequency and linewidth of selected phonon modes in YBCO. We compare the results with experiment, and discuss the features which differ from the single-layer case.

PACS numbers: 74.20.Mn, 74.25.Ha, 74.25.Kc

1 Introduction

Bilayer models for high- T_c superconductors have become popular recently, not only due to the obvious feature that the best-characterised material has a bilayer structural unit, but also since the suggestion by Millis and Monien [1] that the single- and bilayer systems may have fundamentally different magnetic excitations. These differences may include the existence or absence of the spin gap [3, 2], which has become a crucial experimental feature and theoretical benchmark.

In this work we will pursue an approach based on the t - J model [4, 5] analysed by the slave-boson decomposition [6, 7, 8, 9]. In the mean-field approximation, this approach has been studied intensively [10, 11, 12, 13, 14, 15, 16, 17] with a view to describing the magnetic properties of the high- T_c materials. As an example of such investigations, the extended t - J model of a single CuO_2 layer, has been shown [11, 17] to give a good account at the RPA level of many features of the spin excitations in the monolayer $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ (LSCO) system, and some of those in the bilayer $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ (YBCO) case. These authors considered the (in)commensuration of magnetic fluctuations, the frequency dependence of the dynamic susceptibility and the different temperature dependences of the shift and rate of nuclear magnetic resonance between high- and low-doping regions. Such models also contain a consistent description of the “spin-gap” phenomenon [2], first noted by Yasuoka [3]. Taking fluctuations about the mean-field solution into account by a gauge-field approach [18], it is possible to gain an understanding of various transport properties, including the temperature dependence of the resistivity, thermopower and Hall coefficient. Again within the same framework, it has recently been shown [19] that one may obtain a good account of several lattice-related features of the electronic system, such as phonon anomalies and an isotope effect, by considering the coupling which arises naturally between phonon modes of the layer and the spin sector.

In view of this degree of correspondence to experiment, we consider the extended t - J model to be one of the leading candidates for a framework in which to construct a coherent understanding of the many and complex features of the high- T_c problem, and thus that the detailed computation of physical quantities within it is a valuable exercise. Here we wish to apply the ideas of the single-layer studies [17] and [19] to a consistent bilayer model, and thus to elucidate the successes and limitations of the mean-field approach to spin-dependent, microscopic properties. We draw attention to a brief study [13] which established that the bilayer dynamic susceptibility has the experimentally observed q_z periodicity. Ubbens and Lee [16] have also considered static magnetic properties in a bilayer, nearest-neighbour t - J model, including gauge-field effects, and argue for an anisotropic (or “extended”) s -wave pairing state of coupled intra-and interplane singlet order, as well as for enhanced spin-gap formation. Similar results for the spin gap were obtained in Ref. [20].

The question of the gap symmetry has remained a topic of much debate: while it is known in the single-layer model that the low-energy state is d -symmetric [7, 8], a variety of reports find somewhat different behaviour in the presence of coupling to a second layer. In addition to the result of Ref. [16], the weak-coupling antiferromagnetic

spin fluctuation theory, which favours d -symmetry in a single plane, has also been shown to favour extended s -symmetry in a bilayer [21]. A recent numerical analysis by Eder *et al.* [22] gives the evolution of the in-plane d -wave state as interplane coupling is increased, and demonstrates in a small cluster that there is an abrupt crossover to a predominantly interplane ordered state at some value of the coupling. We will discuss these results in comparison with our own in section 2.

The bilayer spin system has also been studied extensively by Millis and coworkers [1, 20, 23, 24], with the aim of elucidating the quantum nature of the ground state and explaining the contrasting NMR results in single- and bilayer materials. From spin-wave and scaling theories [23] it has been shown that interlayer coupling acts to provide a strong enhancement of the spin gap state (by suppression of competing magnetic order), which provides an explanation of the differences without the requirement that the single- and double-layer systems have completely different ground states. These considerations have been cast in a more microscopic form to describe electrons on a bilayer with a weak interplane antiferromagnetic interaction [20]: the system may vary between a Fermi liquid state, which is proposed to describe the doping regime of materials with no spin gap, and a spin liquid picture of fermions interacting via a gauge field, which shows spin-gap features.

The structure of this paper is as follows. In section 2 we consider the mean-field equations of the bilayer problem, deduce the nature of the gap and solve for BSCCO- and YBCO-like systems. In section 3 we calculate the dynamic susceptibility for both types of system, and illustrate the dependence of the results on the wavevector component q_z . In section 4 we compute the superconductive anomalies in c -axis phonon modes of in-plane oxygen atoms for the YBCO system, showing the contrast between modes of even (g) and odd (u) symmetry. Section 5 contains a summary and concluding discussion.

2 Mean-Field Solution

We consider a three-dimensional system consisting of a periodic but uncoupled stack of coupled CuO_2 bilayers, in which the intra- and interplane hopping (t) and superexchange (J) interactions are as shown in Fig. 1. Denoting the c -axis dimension of the unit cell by c and the separation of the planes of the bilayer by d , the ratio $r_c = d/c$ will appear in a phase factor arising from the bilayer spacing. The Hamiltonian for a system of N_z bilayer units is

$$H = \sum_{n=1}^{N_z} \left\{ \sum_{l=1,2} \left[- \sum_{ij\sigma} t_{ij} a_{i\sigma}^{l\dagger} a_{j\sigma}^l + \sum_{\langle ij \rangle} J_{ij} \mathbf{S}_i^l \cdot \mathbf{S}_j^l \right] - \sum_{ij\sigma} t_{\perp ij} \left(a_{i\sigma}^{1\dagger} a_{j\sigma}^2 + a_{i\sigma}^{2\dagger} a_{j\sigma}^1 \right) + \sum_i J_{\perp} \mathbf{S}_i^1 \cdot \mathbf{S}_i^2 \right\}, \quad (1)$$

where l is the layer index in each unit cell. The first line contains the t - J model of a single CuO_2 layer, where t_{ij} denotes the extended transfer integrals, J_{ij} the superex-

change interaction, which is assumed to be finite only between nearest neighbours, and the Hilbert space excludes double occupancy of the quasiparticles $a_{i\sigma}$. The second line contains the coupling between the layers, which may be by both hopping and superexchange. Within the plane, the nearest-neighbour hopping term is taken to be $t = 4J$, while second- and third-neighbour terms will be chosen below in order to match the Fermi surface shape of the physical systems [11]. We will consider initially a range of values for the interlayer coupling parameters $t_{\perp ij}$ and $J_{\perp}$.

The assumption of coherent, single-particle hopping processes between the planes of the bilayer is counter to the interlayer pair tunnelling theories [25], which demands that this vanish by orthogonality. A clear experimental indication of whether interlayer transfer is coherent or incoherent should be given by photoemission experiments, which will see either two Fermi surfaces or one. At the moment there is some controversy, and inconsistency, surrounding the interpretation of results obtained for both BSCCO [26, 27] and YBCO [28, 29] systems, part of which centres on the issue of the number of true band-crossings observed where the bonding and antibonding bands of the coherently-coupled bilayer are expected to be far apart in the Brillouin zone. Here we avoid further discussion of this topic and proceed to elucidate the properties of the bilayer system with coherent hopping, but will refer below to the 2-band interpretations of the experimental data.

In the slave-boson decomposition, the operator a_{js} is represented as $a_{js} = f_{js}b_j^{\dagger}$, where f_{js} is a fermion (spinon) carrying the spin degrees of freedom $\mathbf{S}_j$, and b_j a bosonic holon carrying the charge. In the mean-field approximation, the spin and charge behave independently, and all 4-operator terms may be written expressions quadratic in the operators with the mean-field order parameters (finite expectation values of possible 2-operator combinations) as coefficient, while terms quadratic in these order parameters are also generated. Here we take as possible in-plane order parameters

- i) $\chi_{i,i+\tau} = \sum_{\sigma} \langle f_{i\sigma}^{\dagger} f_{i+\tau\sigma}^{\dagger} \rangle$: uniform RVB of spinons
- ii) $\Delta_{i,i+\tau} = \frac{1}{\sqrt{2}} \langle f_{i\uparrow}^{\dagger} f_{i+\tau\downarrow}^{\dagger} - f_{i\downarrow}^{\dagger} f_{i+\tau\uparrow}^{\dagger} \rangle$: singlet RVB
- iii) $B_{i,i+\tau} = \langle b_{i\sigma}^{\dagger} b_{i+\tau\sigma}^{\dagger} \rangle$: uniform RVB of holons

and as interplane order parameters the combinations

- i) $\chi'_{i,j} = \sum_{\sigma} \langle f_{i\sigma}^{\dagger} f_{j\sigma}^{\dagger} \rangle = \sum_{\sigma} \langle f_{i\sigma}^{2\dagger} f_{j\sigma}^1 \rangle$: interplane uniform RVB of spinons
- ii) $\Delta'_{i,j} = \frac{1}{\sqrt{2}} \langle f_{i\uparrow}^1 f_{j\downarrow}^2 - f_{i\downarrow}^1 f_{j\uparrow}^2 \rangle = \frac{1}{\sqrt{2}} \langle f_{i\uparrow}^2 f_{j\downarrow}^1 - f_{i\downarrow}^2 f_{j\uparrow}^1 \rangle$: interplane singlet
- iii) $B'_{i,j} = \langle b_{i\sigma}^1 b_{j\sigma}^{2\dagger} \rangle = \langle b_{i\sigma}^2 b_{j\sigma}^{1\dagger} \rangle$: interplane uniform RVB of holons.

For historical reasons [30], the terminology “uniform RVB” is taken to signify the hopping order parameter for spin or charge, whose finite value signifies diagonal coherence of the motion. As will be shown below, for the spin order parameters it is sufficient to consider only those between nearest-neighbour sites, so that $\tau = \pm \mathbf{x}$ or $\pm \mathbf{y}$ for χ and Δ , and $\tau = \pm r_c \mathbf{z}$ for χ' and Δ' . In the single-layer problem, the free energy is minimised when $\chi_{i,i+\tau}$ is uniform, but when $\Delta_{i,i\pm x} = -\Delta_{i,i\pm y}$ [7, 8], so that in reciprocal space the gap function Δ_k has d -wave symmetry.

With these definitions, the mean-field approximation to the Hamiltonian (1) in reciprocal space can be represented as

$$H = H_0 + \sum_k \mathbf{b}_k^\dagger \mathbf{M}_{bk} \mathbf{b}_k + \sum_k \mathbf{f}_k^\dagger \mathbf{M}_{fk} \mathbf{f}_k, \quad (2)$$

where H_0 contains terms quadratic in the order parameters, and $\mathbf{b}_k^\dagger = (b_k^{1\dagger}, b_k^{2\dagger})$ and $\mathbf{f}_k^\dagger = (f_{k\uparrow}^{1\dagger}, f_{-k\downarrow}^1, f_{k\uparrow}^{2\dagger}, f_{-k\downarrow}^2)$ are the boson and fermion state vectors. The boson and fermion matrices are given by

$$\mathbf{M}_{bk} = \begin{pmatrix} -\omega_k & -\frac{1}{2}t_\perp^0 \chi_k'^* \\ -\frac{1}{2}t_\perp^0 \chi_k' & -\omega_k \end{pmatrix}, \quad (3)$$

and

$$\mathbf{M}_{fk} = \begin{pmatrix} -\xi_k & \Delta_k & -M_{-k} & D_{-k} \\ \Delta_k & \xi_k & D_k^* & M_k^* \\ -M_{-k}^* & D_k & -\xi_k & \Delta_k \\ D_{-k}^* & M_k & \Delta_k & \xi_k \end{pmatrix} \quad (4)$$

with the following definitions. The band energies for the quasiparticles are $\xi_k = -[t_{fk} + t'_{fk} + t''_{fk} + \frac{3}{4}J\chi_k] - \mu$ and $\omega_k = -[t_{bk} + t'_{bk} + t''_{bk}] - \lambda$, in which the in-plane hopping terms are $t_{fk}^\alpha = \sum_{k'} t_{k'-k}^\alpha \langle b_{k'}^\dagger b_{k'} \rangle$ and $t_{bk}^\alpha = \sum_{k'\sigma} t_{k'-k}^\alpha \langle f_{k'\sigma}^\dagger f_{k'\sigma} \rangle$, with $t_k^\alpha = \sum_{\tau\alpha} t^\alpha e^{i\mathbf{k}\cdot\boldsymbol{\tau}}$, and μ and λ are fermion and boson chemical potentials. The d -symmetric gap is given by $\Delta_k = \frac{3\sqrt{2}}{4}J\Delta(\cos k_x - \cos k_y)$. The effective uniform interplane order parameter is $M_k = t_{\perp k} B_k' + \frac{3}{8}J_\perp \chi_k'$, where the k dependence of $t_{\perp k}$ will be discussed in detail below, and the interplane gap parameter is $D_k = \frac{3\sqrt{2}}{8}J_\perp \Delta_k'$.

The 4×4 unitary matrix which diagonalises the Hamiltonian of the fermionic part may be written as

$$\mathbf{T}_k^{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} \cos \theta_k^+ & \sin \theta_k^+ & -e^{-i\phi} \cos \theta_k^- & -e^{-i\phi} \sin \theta_k^- \\ -\sin \theta_k^+ & \cosh \theta_k^+ & e^{-i\phi} \sin \theta_k^- & -e^{-i\phi} \cos \theta_k^- \\ e^{i\phi} \cos \theta_k^+ & e^{i\phi} \sin \theta_k^+ & \cos \theta_k^- & \sin \theta_k^- \\ -e^{i\phi} \sin \theta_k^+ & e^{i\phi} \cos \theta_k^+ & -\sin \theta_k^- & \cosh \theta_k^- \end{pmatrix}, \quad (5)$$

where

$$\cos 2\theta_k^\pm = \frac{\xi_k \pm |M_k|}{E_k^\pm} \quad \text{and} \quad \sin 2\theta_k^\pm = \frac{\Delta_k \pm |D_k|}{E_k^\pm}. \quad (6)$$

The phase $\phi = -q_z r_c$ is chosen to cancel the q_z -dependent phase of the interplane order parameters, and will be the same for both M_k and D_k ; the modulus in $|Q_k|$ thus denotes the absence of this phase, but does not prevent the order parameter from being negative. It is the transformation (5) which will be used extensively in the sections to follow, when the boson operators $\mathbf{f}_k$ are replaced in the physical quantities to be computed by the quasiparticle operators $\vec{\gamma}_k = \mathbf{T}_k \mathbf{f}_k$.

The fermion part of the Hamiltonian then takes the diagonal form

$$\widetilde{\mathbf{M}}_{fk} = \begin{pmatrix} -E_k^+ & 0 & 0 & 0 \\ 0 & E_k^+ & 0 & 0 \\ 0 & 0 & -E_k^- & 0 \\ 0 & 0 & 0 & E_k^- \end{pmatrix}, \quad (7)$$

in which the eigenvalues are given by the combinations

$$E_k^\pm = \left[(\xi_k \pm |M_k|)^2 + (\Delta_k \pm |D_k|)^2 \right]^{1/2}. \quad (8)$$

E_k^- corresponds to the bonding band of the bilayer system, and E_k^+ to the antibonding band.

The 2×2 bosonic matrix (3) is trivially diagonalised by the transformation

$$b_k^1 = \frac{1}{\sqrt{2}} (\beta_k^1 - e^{-i\phi} \beta_k^2) \quad (9)$$

$$b_k^2 = \frac{1}{\sqrt{2}} (e^{i\phi} \beta_k^1 - \beta_k^2) \quad (10)$$

to new quasihole operators β_k , in terms of which the diagonal boson matrix is

$$\widetilde{\mathbf{M}}_{bk} = \begin{pmatrix} -\tilde{\omega}_k^+ & 0 \\ 0 & -\tilde{\omega}_k^- \end{pmatrix}, \quad (11)$$

where the eigenenergies are given by

$$\tilde{\omega}_k^\pm = \omega_k \pm \frac{1}{2} t_\perp^0 |\chi'_k|. \quad (12)$$

The system is required to satisfy eight mean-field equations, three for the in-plane order parameters, three for the interplane parameters and two for the carrier number δ . These equations may be obtained by minimising the free energy with respect to each of the parameters in turn, or by substituting the quasiparticle operators into the reciprocal-space definitions of the order parameters and transforming back to real space. One obtains for the in-plane parameters

$$\chi_{i,i+\tau} = -\frac{1}{N} \sum_k e^{-i\mathbf{k}\cdot\tau} \left[\cos 2\theta_k^+ \tanh \frac{1}{2} \beta E_k^+ + \cos 2\theta_k^- \tanh \frac{1}{2} \beta E_k^- \right], \quad (13)$$

$$\Delta_{i,i+\tau} = -\frac{1}{\sqrt{2}N} \sum_k e^{-i\mathbf{k}\cdot\tau} \left[\sin 2\theta_k^+ \tanh \frac{1}{2} \beta E_k^+ + \sin 2\theta_k^- \tanh \frac{1}{2} \beta E_k^- \right] \quad (14)$$

and

$$B_{i,i+\tau} = \frac{1}{N} \sum_k e^{-i\mathbf{k}\cdot\tau} \left[n(\tilde{\omega}_k^+) + n(\tilde{\omega}_k^-) \right], \quad (15)$$

for the interplane parameters

$$\chi'_{i,i+\tau} = -\frac{1}{N} \sum_k \left[\cos 2\theta_k^+ \tanh \frac{1}{2} \beta E_k^+ - \cos 2\theta_k^- \tanh \frac{1}{2} \beta E_k^- \right], \quad (16)$$

$$\Delta'_{i,i+\tau} = -\frac{1}{\sqrt{2N}} \sum_k \left[\sin 2\theta_k^+ \tanh \frac{1}{2}\beta E_k^+ - \sin 2\theta_k^- \tanh \frac{1}{2}\beta E_k^- \right] \quad (17)$$

and

$$B'_{i,i+\tau} = \frac{1}{N} \sum_k e^{-i\mathbf{k}\cdot\tau} \left[n(\tilde{\omega}_k^+) - n(\tilde{\omega}_k^-) \right], \quad (18)$$

and for the carrier number, which must be equal to the number of bosons, and by conservation must add to the number of fermions to give unity,

$$\delta = \frac{1}{2N} \sum_k \left[n(\tilde{\omega}_k^+) + n(\tilde{\omega}_k^-) \right] \quad (19)$$

from the boson part and

$$\delta = \frac{1}{2N} \sum_k \left[\cos 2\theta_k^+ \tanh \frac{1}{2}\beta E_k^+ + \cos 2\theta_k^- \tanh \frac{1}{2}\beta E_k^- \right] \quad (20)$$

from the fermion part. In (13 - 20), the sine and cosine factors are given by (6) and $n(\omega_k)$ is the Bose occupation function.

In this type of model, the bosonic holons occupy only a very small region of reciprocal space close to $\mathbf{k} = 0$, and in all respects are effectively condensed at all reasonable temperatures. Thus to a very good approximation one may take $B = \delta$ in (15) and $B' = \delta \text{sgn}(\chi')$ in (18), so that the bosonic equations need not be considered further. It is this approximation which allows us to consider only nearest-neighbour spin order parameters in a model with only nearest-neighbour superexchange interactions. Thus the problem reduces to a five-parameter one in the fermionic degrees of freedom, with the mean-field equations (13), (14), (16), (17) and (20).

Following Tanamoto *et al.* [11, 17], we concentrate on a doping level close to the value where the system is close to reproducing the properties of optimally-doped materials, and choose $\delta = 0.2$. Here we expect a solution with finite intra- and interplane uniform order parameters over a considerable range of temperatures, and this is borne out by calculations. In the low-temperature regime we wish to investigate the nature of the transition to a state with singlet order, and the possible coexistence or competition of the intra- and interplane ordered states which may exist here. The idea of a singlet state with coupled order parameters has been studied in one version of the current model by Ubbens and Lee [16], and proposed by Kuboki and Lee [31] as a candidate which may explain recent observations of the anisotropic gap in BSCCO materials by photoemission [27]. A very recent numerical study [22] has investigated the same issue in small, bilayer clusters, finding a sharp crossover from predominantly in-plane to predominantly interplane singlet order at a value of the interlayer superexchange parameter $J_\perp = xJ$, where x is of order unity, but may under certain circumstances be quite small. We will compare our results with those of Ref. [22] in more detail below.

In a single-layer system, the gap parameter Δ may have an arbitrary phase θ , from which no physical consequences arise. In a bilayer, the phases of the parameters on each layer may differ, thus appearing as $\Delta_k e^{i\theta_1}$ and $\Delta_k e^{i\theta_2}$. When the layers are coupled, there

may in general be an additional term in the Hamiltonian of the form $H = -\epsilon \cos(\theta_1 - \theta_2)$, where the energy scale of the phase coupling parameter is $\epsilon \sim t_\perp^0{}^2/W$, in which W is the bandwidth of the in-plane dispersion. This term may couple to an external magnetic field, and gives rise to a variety of Josephson coupling phenomena, some of which are integral to the qualitatively new predictions of the interlayer pair tunnelling theory [25]. In this work, we assume that the minimum energy state is maintained, so that the phases in the two layers remain equal, and both gap functions can be taken simply to be Δ_k , as given below (4).

In order to make the following study more directly applicable to the physical systems, we appeal to the results of Andersen *et al.* [32] for some additional detail about the interplane hopping term. From detailed bandstructure studies on the YBCO system, these authors estimate the magnitude to be $t_\perp^0 \simeq 0.15t$, where in our choice of units the maximal band splitting will be $8t_\perp^0$. The k dependence resulting from the combination of the interplane term with the extended in-plane terms is contained consistently in the extended interplane hopping parameters $t_\perp$, $t'_\perp$ and $t''_\perp$ (Fig. 1), and is such that $t_\perp(\mathbf{k})$ is small along the ΓM line in the Brillouin zone. This has been parameterised in the form

$$t_\perp(\mathbf{k}) = t_\perp^0 (\cos k_x - \cos k_y)^2, \quad (21)$$

which corresponds to the special case where the bands are degenerate along ΓM , and results from a real-space distribution of the interplane hopping integrals given by $t_\perp = 0$ [32], $t'_\perp = -\frac{1}{2}t_\perp^0$ and $t''_\perp = \frac{1}{4}t_\perp^0$. The interlayer spin exchange is taken to be nearest-neighbour only, and to have a small value $J_\perp \sim 0.1J$, in line with the majority of experimental [33, 34, 35] and theoretical [32] indications. We note, however, that recent infrared transmission and reflectometry measurements [36] suggest a considerably larger value of this parameter in the insulating YBCO system, and so we do not discount this possibility.

To investigate the nature of the spin-ordered state, we consider the linearised gap equations (14) and (17) close to the their transition temperature. In their most general form these contain the 3 equations

$$\begin{aligned} \Delta_x &= \frac{3}{4} \sum_k \cos k_x \left(\frac{\Delta_x \cos k_x + \Delta_y \cos k_y + \frac{1}{2} J_\perp \Delta_\perp}{2\xi_k^+} \tanh \frac{1}{2} \beta \xi_k^+ \right. \\ &\quad \left. + \frac{\Delta_x \cos k_x + \Delta_y \cos k_y - \frac{1}{2} J_\perp \Delta_\perp}{2\xi_k^-} \tanh \frac{1}{2} \beta \xi_k^- \right) \\ \Delta_y &= \frac{3}{4} \sum_k \cos k_y \left(\frac{\Delta_x \cos k_x + \Delta_y \cos k_y + \frac{1}{2} J_\perp \Delta_\perp}{2\xi_k^+} \tanh \frac{1}{2} \beta \xi_k^+ \right. \\ &\quad \left. + \frac{\Delta_x \cos k_x + \Delta_y \cos k_y - \frac{1}{2} J_\perp \Delta_\perp}{2\xi_k^-} \tanh \frac{1}{2} \beta \xi_k^- \right) \\ \Delta_\perp &= \frac{3}{4} \sum_k \left(\frac{\Delta_x \cos k_x + \Delta_y \cos k_y + \frac{1}{2} J_\perp \Delta_\perp}{2\xi_k^+} \tanh \frac{1}{2} \beta \xi_k^+ \right. \end{aligned} \quad (22)$$

$$- \frac{\Delta_x \cos k_x + \Delta_y \cos k_y - \frac{1}{2} J_\perp \Delta_\perp}{2\xi_k^-} \tanh \frac{1}{2} \beta \xi_k^- \Big),$$

where $\xi_k^\pm = \xi_k \pm |M_k|$ and $\Delta_\perp$ denotes Δ' . This set of equations can be cast in the schematic matrix form

$$\begin{pmatrix} I_1^{xx} + I_2^{xx} - 1 & I_1^{xy} + I_2^{xy} & I_1^x - I_2^x \\ I_1^{xy} + I_2^{xy} & I_1^{yy} + I_2^{yy} - 1 & I_1^y - I_2^y \\ I_1^x - I_2^x & I_1^y - I_2^y & I_1 + I_2 - \frac{2}{J_\perp} \end{pmatrix} \begin{pmatrix} \Delta_x \\ \Delta_y \\ \frac{1}{2} J_\perp \Delta_\perp \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad (23)$$

where $I_\alpha = \langle 1 \rangle_\alpha$, $I_\alpha^i = \langle \cos k_i \rangle_\alpha$ and $I_\alpha^{ij} = \langle \cos k_i \cos k_j \rangle_\alpha$, in which the angled brackets $\langle \rangle_\alpha$ denote integration over the Brillouin zone of the energy denominator and thermal function of the energy branch α . A coupled transition of all 3 order parameters may occur where the determinant of this matrix is zero. The determinantal equation may be represented simply by

$$\begin{vmatrix} a & b & c \\ b & a & c \\ c & c & d \end{vmatrix} = (a - b) [(a + b)d - 2c^2] = 0. \quad (24)$$

The in-plane d -symmetric solution is sought by writing the eigenvector $(\Delta_x, \Delta_y, \Delta_\perp)$ as $(\Delta, -\Delta, \Delta_\perp)$, where Δ is real, and it is clear that the only possible solution is $\Delta_\perp = 0$. This is the root $a - b = 0$ of (24) above. Thus in the mean-field framework, d -wave singlet pairing in the plane is thus found to exclude any interplane singlet order at the same onset temperature $T = T_{RVB}$. This is the most important qualitative result of the current analysis, as it has profound implications for the nature of the singlet spin-ordered state which may be realised in this type of model. The in-plane s -wave solution $(\Delta, \Delta, \Delta_\perp)$ can be found to exist with $\frac{\Delta_\perp}{\Delta} = -\frac{2c}{d}$, which is the root $[(a + b)d - 2c^2] = 0$ in (24) above. This is a coupled solution with in-plane s -wave symmetry, which for small $J_\perp$ admits some admixed interplane singlet order in an amount proportional to the band splitting, and at large $J_\perp$ represents a predominantly interplane singlet-ordered state. We have not been able to find a solution to the system of mean-field equations for the in-plane s -wave ordered state for any values of the doping δ or interplane superexchange interaction $J_\perp$ in the regime of physical interest, and so do not consider this state further here. We will have some further comment on the interplane ordered state below, in the context of its possible coexistence with in-plane d -wave singlet pairing at temperatures away from T_{RVB} .

Returning to consider the case of a d -symmetric in-plane singlet order parameter in more detail, we note first that if the planes are decoupled ($t_\perp^0 = 0$), $\xi_k^+ = \xi_k^-$ and the equations for (Δ_x, Δ_y) and $\Delta_\perp$ decouple. Thus there would be two separate transitions for the total order parameter if interplane singlet ordering were not excluded as in the coupled case above. To investigate the possibility of coexisting order parameters in some temperature regime of the system, one may consider the in-plane d -symmetric ordered state and seek a second transition to finite interplane order at a lower temperature.

In the absence of interplane singlet order, this state is described by the reduced system of 4 equations ((13), (14), (16) and (20)), which can be solved readily to yield a self-consistent set of solutions for the order parameters as functions of temperature. In Fig. 2 we show the transition temperature T_{RVB} , representing the onset of in-plane singlet order in such a system, as a function of the interplane hopping parameter $t_{\perp}^0$. T_{RVB} is little affected by small interplane hopping, but loses most of its value as $t_{\perp}^0$ (21) approaches J . These results were obtained with a fixed, small interplane spin interaction $J_{\perp} = 0.1J$ at all values of $t_{\perp}^0$.

Returning to the system of 5 equations, this can be studied below T_{RVB} at several values of $t_{\perp}^0$ (Fig. 2) to seek a transition to non-zero $\Delta_{\perp}$ (17). We find that this possibility exists only for values of $0.8J < J_{\perp} < 1.2J$: for smaller values the equation cannot be satisfied because the coefficient $J_{\perp}$ of the left-hand side is too small to compete with the band splitting, and for larger ones it is likely that the assumption of the interplane singlet state being subsidiary to the in-plane state is no longer valid [22]. The range of coexistence is quite insensitive to the value of $t_{\perp}^0$, a feature which we believe is due to the line nodes in $t_{\perp k}$ (21). This result is in agreement with a recent study by Eder *et al.* [22], who investigated spin correlation functions in 16- and 20-site bilayer clusters by exact diagonalisation. These authors find for a doped system a very abrupt crossover, from a state with predominantly in-plane d -wave spin correlation to one of “standing singlets” formed between nearest-neighbour spins in each layer, as a function of increasing interplane superexchange $J_{\perp}$. The crossover occurs at a value $J_{\perp} = xJ$, in which x varies over a parameter-dependent range of values below unity. That interplane singlet order is not compatible with in-plane d -symmetric singlet pairing was also found by Liechtenstein *et al.* [21] in a bilayer antiferromagnetic spin fluctuation model, which was shown to favour the existence of the anisotropic s -wave state.

These results raise the interesting question of how the d -symmetric in-plane singlet order in an isolated plane may evolve as interplane hopping and superexchange increase. One alternative to the admixture of real order parameters considered above, which has been shown not to evolve below some significant threshold value of the coupling, is the development of imaginary components of the order parameter. If the order parameter has the form $\Delta_x = \Delta_1 + i\Delta_2$, $\Delta_y = -\Delta_1 + i\Delta_2$, one may formulate a system of equations analogous to (22) above, in which the real part is the single-layer equation for in-plane d -order and the imaginary part a coupled equation for Δ_2 and a purely imaginary interplane component $i\Delta_{\perp}$. This has been termed the $d + is$ state, and would give a completely nodeless gap because the order parameter has a finite imaginary part where the real part vanishes. However, the real and imaginary parts are decoupled, so there is no reason to expect the two transitions to occur simultaneously. A second transition below T_{RVB} to a nodeless state has been proposed as the explanation for a feature in the NMR spin-echo relaxation rate T_{2G}^{-1} [37], and so this state is of both experimental and theoretical interest. We have sought a transition to this state in the same way as described above, and again found that it did not exist within the present framework below values of $J_{\perp} \sim J$.

A different proposition for the existence of the coupled singlet state is given by the argument [16, 20] that fluctuations of the gauge-field in a bilayer system are considerably more effective in suppressing in-plane d -wave order than interplane s -wave order, and thus act to allow the latter to survive in the presence of the former. The same authors also argue for enhancement of the interlayer coupling $J_{\perp}(\mathbf{q})$, to values which may fall in the coupled regime, close to the antiferromagnetic wavevector (π, π) . Finally, if the system is orthorhombic, *i.e.* $t_x \neq t_y$ and $J_x \neq J_y$ in the plane, one may show [31, 38] that the possibility of a coupled transition does exist, and thus that a mixed “ $d + s$ ” or “ $d + is$ ” state of coexisting intra- and interplane order parameters is justifiable in the physical parameter regime. However, there is as yet no definitive evidence that the BSCCO system, whose photoemission spectra have been interpreted to suggest a $d + s$ state [31], shows any orthorhombicity of a type which would couple appropriately to both gap parameters.

The range of $J_{\perp}$ for which a mixed state of in- and interplane singlet spin order may exist is unrealistically large for the high- T_c superconducting materials, and in addition is unphysical in the context of a model where the superexchange interaction should be the result of a second-order hopping process, with magnitude $J_{\perp} \sim \frac{t_{\perp}^0{}^2}{U}$. In order to investigate the properties to be expected from the model, and to compare them with experiment, we wish to restrict the discussion to the physical parameter regime, which will require considerably smaller values of $J_{\perp}$. We will thus proceed by taking the interplane singlet order parameter to be zero, and to study the 4-parameter problem for the uniform in- and interplane order parameters χ and χ' , the in-plane spin gap parameter Δ and the chemical potential μ , albeit in the presence of a finite interplane spin coupling constant.

Following Andersen *et al.* [32], we now fix the magnitude of the interplane hopping term at $t_{\perp}^0 = 0.15t$. We choose to set the interplane spin coupling to the small value $J_{\perp} = 0.1J$, noting that if in fact this parameter is found by more detailed experiments to have a somewhat larger value, its role in the structure of the current theory is such that it has a negligible effect on the mean-field solutions and Fermi surface shape. In the spirit of the single-layer analysis of Tanamoto and coworkers [11, 17], we wish to choose the extended transfer integrals t' and t'' in such a way that the Fermi surfaces of the bonding and antibonding bands reproduce as many features as possible of the physical bilayer systems of interest, BSCCO and YBCO. As was shown clearly by these authors, the magnetic properties of the system are strongly influenced by the exact shape of the Fermi surface.

As a prototype of the BSCCO system, in Fig. 3(a) we show Fermi surfaces in the first quadrant of reciprocal space for the parameter choice $t = 4J$, $t' = -\frac{1}{8}t$ and $t'' = \frac{1}{6}t$. For the fixed value of $t_{\perp}^0$, this leads to a pair of dispersion surfaces with flat regions around the $(\pi, 0)$ saddle points, the important feature of which is that the “extended saddle point” regions of the bonding band lie below the chemical potential, while those of the antibonding band lie above it. Thus the bonding (outer) band has the open shape characteristic of YBCO-like systems [17], while the antibonding (inner) band has

the closed, LSCO-like shape; we will use the terms “open” and “closed” to refer to the nature of the Fermi surface in the first Brillouin zone, in an electron picture. In addition, the parameters are chosen so that the antibonding band is rather strongly nested, by which is meant that the Fermi surface has significant regions lying almost on a straight line parallel to $q_x + q_y = 0$, so that the same wavevector $\mathbf{q}$ can span many pairs of Fermi surface points; in detail, this surface is slightly concave in relation to the given line. Finally, the bands appear to be degenerate along the direction ΓM in the Brillouin zone, in accordance with the calculations of Ref. [32], and so we retain the form for $t_{\perp k}$ (21) proposed by these authors. These are the features of the published interpretation of the photoemission data from BSCCO in Ref. [26], and so we take this parameter set to characterise the BSCCO system in the remainder of this study.

To describe the YBCO system, in Fig. 3(b) we show Fermi surfaces in the first quadrant for the parameter choice $t = 4J$, $t' = -\frac{1}{5}t$ and $t'' = \frac{1}{4}t$. In this case the parameters are such that the extended saddle point regions of both bands are below the chemical potential, so that both have open Fermi surfaces. However, the antibonding band is very flat in these regions, and the Fermi surface shows a rather narrow “neck”, so is close to becoming closed on small changes of the parameters. We emphasise in passing that these strong alterations in Fermi surface shape are a consequence of very small changes in the parameters t' and t'' , because of the flat nature of the saddle-point region. Here we require that the bands are not fully degenerate in the ΓM direction, and so alter the form of the interplane term to $t_{\perp k} = \frac{2}{3}[(\cos k_x - \cos k_y)^2 + 2]$. This alteration corresponds to the case where the interplane hopping integrals are given by $t_{\perp} = 0$, $t'_{\perp} = -\frac{1}{3}t_{\perp}^0$ and $t''_{\perp} = \frac{1}{6}t_{\perp}^0$, or in other words that the interplane hopping is less extended in this case. The function in reciprocal space remains minimal along the ΓM . The fitted features are those of the YBCO system, as calculated from first principles in Ref. [32], and are fully consistent with results from the most accurate photoemission studies of YBCO 124 [28] and 123 [29] to date. Henceforth we take this parameter set to give the prototypical YBCO system for analysis of the physical properties. We believe that the relatively large value of $t'' = \frac{1}{4}t$ is reasonable, as it remains consistent with the expectation from microscopic models for the transfer of extended quasiparticles in the CuO_2 plane [39]. The Fermi surfaces shown in Fig. 3 are independent of the perpendicular wavevector k_z , which does not appear in the quasiparticle dispersion (8) due to the assumption of no coupling between unit cells. In the slave-boson decomposition at the mean-field level, because the holons are concentrated at $q = 0$, calculated photoemission spectra are exactly characteristic of the spinon Fermi surfaces (Fig. 3), and contain no alterations reflecting any effects of the charge degrees of freedom on the spin.

With the above choices one may solve the mean-field equations of the 4-parameter fermion system, to obtain all of the parameters as functions of temperature. In Fig. 4 we show the form of the in-plane singlet spin order parameter $\Delta(T)$ for both BSCCO-like (Fig. 4(a)) and YBCO-like (Fig. 4(b)) systems. Δ shows a BCS-like second-order transition, and the transition temperature T_{RVB} sets the characteristic scale for spin-related properties in this type of model. We turn now to the computation of two such

properties, the dynamic susceptibility and the phonon anomalies.

3 Dynamic Susceptibility

The dynamic susceptibility $\chi(\mathbf{q}, q_z, \omega)$ contains all information about the spin response of a magnetic material, at all frequencies and wavevectors. Its zero-frequency limit is sampled at certain wavevectors by NMR experiments, while the full dynamical quantity is measured by inelastic neutron scattering, as discussed in more detail from the viewpoint of this type of model in Ref. [17]. Here we will compute the dynamic susceptibility in the random-phase approximation (RPA), one method by which spin fluctuation enhancement may be taken into account, and illustrate the results as scans of in-plane wavevector $\mathbf{q}$ or frequency ω with the other variable held fixed. We will show also the variation of the susceptibility with the wavevector component q_z , and explain the results in terms of intra- and interband scattering processes.

In a bilayer system, the imaginary part of the bare, retarded susceptibility $\chi_0''(\mathbf{q}, q_z, \omega)$ is a 2×2 matrix which can be calculated by analytic continuation of the associated thermal function

$$\chi_0(\mathbf{q}, q_z, i\omega_n) = \begin{pmatrix} \langle \mathbf{S}^1 \cdot \mathbf{S}^1 \rangle & \langle \mathbf{S}^1 \cdot \mathbf{S}^2 \rangle \\ \langle \mathbf{S}^2 \cdot \mathbf{S}^1 \rangle & \langle \mathbf{S}^2 \cdot \mathbf{S}^2 \rangle \end{pmatrix}, \quad (25)$$

in which $\langle \mathbf{S}^l \cdot \mathbf{S}^m \rangle$ denotes $\langle \mathbf{S}^l(\mathbf{q}, q_z, i\omega_n) \cdot \mathbf{S}^m(\mathbf{q}, q_z, i\omega_n) \rangle$. The spin operator $\mathbf{S}$ may be written in terms of the operators $\vec{f}_k$, transformed to the quasiparticle operators $\vec{\gamma}_k$ and the resulting expressions evaluated in the diagonal basis. Omitting details of this lengthy procedure, the bare susceptibility assumes the form

$$\chi_0(\mathbf{q}, q_z, \omega) = \begin{pmatrix} \chi_+(\mathbf{q}, \omega) & e^{iq_z r_c} \chi_-(\mathbf{q}, \omega) \\ e^{-iq_z r_c} \chi_-(\mathbf{q}, \omega) & \chi_+(\mathbf{q}, \omega) \end{pmatrix}, \quad (26)$$

where the diagonal and off-diagonal parts are quantities whose imaginary parts have the form

$$\begin{aligned} \chi_{\pm}''(\mathbf{q}, \omega) = & \sum_{\alpha\beta} \frac{\pi}{8} A^{\pm} \sum_k \left[C_{k,q}^+ T_{k,q}^+ \left[\delta(\omega - E_k^{\beta} + E_{k+q}^{\alpha}) - \delta(\omega + E_k^{\beta} - E_{k+q}^{\alpha}) \right] \right. \\ & \left. + C_{k,q}^- T_{k,q}^- \left[\delta(\omega + E_k^{\beta} + E_{k+q}^{\alpha}) - \delta(\omega - E_k^{\beta} - E_{k+q}^{\alpha}) \right] \right], \end{aligned} \quad (27)$$

and similarly for the real parts $\chi_{\pm}'$ [13]. Here the indices α and β denote the bonding or antibonding bands, $A^+ = 1$ and $A^- = 2\delta_{\alpha\beta} - 1$,

$$C_{k,q}^{\pm} = 1 \pm \cos 2(\theta_{k+q}^{\alpha} - \theta_k^{\beta}) \quad (28)$$

gives the coherence factors and

$$T_{k,q}^{\pm} = f(\pm E_k^{\beta}) - f(E_{k+q}^{\alpha}) \quad (29)$$

gives the thermal factors. Thus each component of χ_0 is a combination of 16 terms, 8 for normal scattering processes and 8 for anomalous ones. All of the contributions are additive in χ_+ , but the interband contributions ($\alpha \neq \beta$) are subtractive in χ_- .

Note that the only dependence on q_z is that contained explicitly in the phase factors of the off-diagonal terms in (26). This phase will appear in the imaginary part as the legitimate coefficient of terms proportional to a δ -function in energy which are responsible for dissipative processes, and is not forbidden by the symmetry of χ'' . On taking the symmetric combinations required to compute any physical quantity, the phases will combine to give a real result.

The RPA susceptibility is a sum of chains of bare polarisation parts connected by the spin interaction $\mathbf{J}(\mathbf{q}, q_z)$, and can be written in the form of a matrix Dyson equation as

$$\chi(q, \omega) = \chi_0(q, \omega) - \chi_0(q, \omega) \mathbf{J}(q) \chi(q, \omega) \quad (30)$$

$$= (\mathbf{1} + \chi_0(q, \omega) \mathbf{J}(q))^{-1} \chi_0(q, \omega), \quad (31)$$

where q denotes $(\mathbf{q}, q_z)$. The matrix form of the spin interaction on the bilayer is

$$\mathbf{J}(\mathbf{q}, q_z) = \begin{pmatrix} J_{\parallel}(\mathbf{q}) & e^{iq_z r_c} J_{\perp}(\mathbf{q}) \\ e^{-iq_z r_c} J_{\perp}(\mathbf{q}) & J_{\parallel}(\mathbf{q}) \end{pmatrix}, \quad (32)$$

in which by the assumption of nearest-neighbour interactions only, one has $J_{\parallel}(\mathbf{q}) = J(\cos q_x + \cos q_y)$ and $J_{\perp}(\mathbf{q}) = J_{\perp}$.

Omitting the functional dependences and superscripts for clarity, the RPA susceptibility from (31) has the form

$$\chi = \frac{1}{|det|} \begin{pmatrix} \chi_+ + J_{\parallel}(\chi_+^2 - \chi_-^2) & e^{i\phi} (\chi_- + J_{\perp}(\chi_+^2 - \chi_-^2)) \\ e^{-i\phi} (\chi_- + J_{\perp}(\chi_+^2 - \chi_-^2)) & \chi_+ + J_{\parallel}(\chi_+^2 - \chi_-^2) \end{pmatrix}, \quad (33)$$

where ϕ denotes $q_z r_c$ and the determinant of the matrix $(\mathbf{1} + \chi_0 \mathbf{J})$ has the separable form

$$\begin{aligned} |det| &= 1 + 2(\chi_+ J_{\parallel} + \chi_- J_{\perp}) + (\chi_+^2 - \chi_-^2)(J_{\parallel}^2 - J_{\perp}^2) \\ &= (1 + (J_{\parallel} + J_{\perp})(\chi_+ + \chi_-))(1 + (J_{\parallel} - J_{\perp})(\chi_+ - \chi_-)). \end{aligned} \quad (34)$$

The measured quantity in an experiment such as the determination of the neutron scattering cross section is the sum of all components of the imaginary part of the susceptibility matrix, *i.e.*

$$S^R(q, \omega) \propto \sum_{\mu\nu} \frac{\chi''_{\mu\nu}(q, \omega)}{1 - e^{-\beta\omega}}, \quad (35)$$

where μ and ν denote the layer index. Thus the diagonal terms in (33) are the effective RPA susceptibility for spins in a single plane of the bilayer, and the off diagonal terms give the interplane contribution, which will be proportional to $\cos q_z r_c$.

By performing the summation and separation, the final form of the cross section is

$$S^R(q, \omega) \propto \text{Im} \left\{ \frac{(\chi_+ + \chi_-) \cos^2 \frac{1}{2} q_z r_c}{1 + (J_{\parallel} + J_{\perp})(\chi_+ + \chi_-)} + \frac{(\chi_+ - \chi_-) \sin^2 \frac{1}{2} q_z r_c}{1 + (J_{\parallel} - J_{\perp})(\chi_+ - \chi_-)} \right\}, \quad (36)$$

a more general version of the result given by Ref. [16]. Returning to the equation (27), one observes that the components may be cast in the form $\chi_{\pm} = \chi_{\alpha\alpha} \pm \chi_{\alpha\beta} \pm \chi_{\beta\alpha} + \chi_{\beta\beta}$, where each of the terms $\chi_{\alpha\alpha'}$ has contributions from 2 normal and 2 anomalous scattering processes, and is either purely intra- ($\alpha = \alpha'$) or interband. The 2 terms in (36) are then either purely intraband ($\chi_+ + \chi_- = 2(\chi_{\alpha\alpha} + \chi_{\beta\beta})$) or interband ($\chi_+ - \chi_- = 2(\chi_{\alpha\beta} + \chi_{\beta\alpha})$) in nature. Thus the wavevector q_z may be used to select which type of scattering process is measured, so that the contributions of each may be compared.

In Figs. 5-7 we give results for the wavevector and frequency dependence of the imaginary part of the dynamic susceptibility $\chi = \sum_{\mu\nu} \chi_{\mu\nu}$ for the BSCCO- and YBCO-like systems, based on the Fermi surface parameters chosen in section 2. Here we will show only results for the system at low temperatures (specifically, $T = 0.1T_{RVB}$), so that it is in the singlet RVB state; the qualitative features of the evolution from the high-temperature state are well illustrated in Ref. [17]. In Fig. 5(a) is shown $\chi''(\mathbf{q}, \omega)$ for the BSCCO system as $\mathbf{q}$ is scanned along the symmetry directions of the Brillouin zone. The frequency is fixed at $\omega = 0.3J$, above the value of the twice the maximal gap Δ_k , and the out-of-plane wavevector at $q_z = 0$ so that only scattering processes within the bonding and antibonding bands may contribute. One observes a significant incommensuration due to the strongly-nested shape of the antibonding band, and a general increase in the scattered intensity for values of $\mathbf{q}$ near $\mathbf{Q} = (\pi, \pi)$, the antiferromagnetic wavevector which spans many points close to the Fermi surface of the bonding band (Fig. 3). This is a combination of the Fermi-surface features documented in Ref. [17] for both closed and open types, which clearly vary little with energy transfer. In Fig. 6(a) is shown the same quantity for $q_z = \pi$ (in “bilayer units”, $r_c = 1$), so that only bonding to antibonding band scattering processes are sampled. Here the scattered intensity is considerably higher, indicating a greater enhancement from the relevant denominator in (36), and is largely commensurate as a result of poor nesting between the two bands. In Fig. 7(a) is shown $\chi''(\mathbf{q}, \omega)$ at fixed wavevector $\mathbf{q} = \mathbf{Q}$ as the frequency is scanned up to values of order J , for 3 choices of q_z and at low temperature. The results for interband scattering are very similar to those obtained in Ref. [17] for the single layer with open Fermi surface, in that there is a strong peak due to states pushed out of a gap ($\Delta_{\alpha}(\mathbf{k}_{\alpha}) + \Delta_{\beta}(\mathbf{k}_{\beta})$) here, with $\mathbf{k}_{\alpha} - \mathbf{k}_{\beta} = (\pi, \pi)$, followed by a broad region of spectral weight up to a shoulder at $\omega \sim J$ which characterises the maximal energy separation of significant numbers of points whose wavevector separation is (π, π) for the energy dispersions of the system. For intraband scattering, the shoulder appears at a similar energy, but the peak feature is missing; this can be understood due to the fact that there is no contribution from processes involving the antibonding band, all of whose spanning wavevectors are smaller than $\mathbf{Q}$, with the result that the denominator in the first term of (36) is far from being small, and so provides very little enhancement.

Fig. 5(b) shows the variation of χ'' with $\mathbf{q}$ in the Brillouin zone at fixed frequency, low temperature and $q_z = 0$ for the YBCO system, where it is clear that the spin excitations are commensurate, and strong over a broad range of $\mathbf{q}$ around (π, π) . This result appears from [17] to be characteristic of the open Fermi surface shape; the only qualitative effect of the higher energy transfer shown here in the singlet RVB state is that one no longer observes the peaks which arise due to scattering processes between the gap nodes. For $q_z = \pi$ (Fig. 6(b)), the peak is again commensurate, as is to be expected as the 2 bands involved both have the open shape, and once again has higher intensity than that from intraband processes. Fig. 7(b) shows that the excitation spectrum as a function of ω once more has a shape similar to that seen in the single-layer case, with the exception of clear evidence of 2 separate contributions to the peaks at each q_z . This is the result of having 2 pairs of Fermi-surface points separated by (π, π) , so that the characteristic gap combinations in each case may differ, and is most pronounced in the $q_z = \pi$ case. The shoulder feature for the YBCO bands occurs at $\omega \sim 1.2J$.

We may conclude by noting that at the RPA level the dynamic susceptibility is very much dominated by Fermi-surface scattering processes, which are strongly favoured over even those involving nearby points, and in this sense the results show little qualitative difference from those of the single layer. The calculated result that interplane contributions are stronger than intraplane ones over most of the energy range (Fig. 7) arises primarily from their smaller enhancing denominator (36), and is consistent with the results of experiment in YBCO [40]. In the absence of coherent interplane hopping [25], a different explanation of this modulation would be required. However, with regard to overall agreement with experimental observation, we do not consider the results given in Figs. 5-7(b) to be a satisfactory description of the measured magnetic response of YBCO materials [41, 42, 43]. Recent experiments show few features in the spin excitation spectrum other than a strong and remarkably narrow peak at 41meV in the superconducting state, although there is evidence for some broad spectral weight below this energy in the normal state. The peak occurs only close to the wavevector $\mathbf{Q}$, and does not seem to disperse significantly with $\mathbf{q}$ or ω , so that the band picture given above, and proposed by many authors, would appear to be a poor candidate for a full explanation. This is in contrast to the spin spectrum of the single-layer LSCO material [17, 44, 45], where the current form of theory gives a good account of the experimental features. In YBCO, the narrow and non-dispersive nature of the 41meV peak suggests the existence of a resonance, and there has been a theoretical proposal for this within the class of tight-binding, strongly-correlated systems in two dimensions [46]. A full investigation of this possibility in the present framework will be the subject of a subsequent publication. Finally, we comment that no measurements of dynamic susceptibility have so far been performed on BSCCO materials, because these have proven extremely difficult to grow as high-quality single crystals of the required size [47].

4 Phonon Anomalies

The coupling between the spin and lattice degrees of freedom in the t - J model has been investigated recently in Ref. [19]. These authors use the modulation of the parameters t_{ij} and J_{ij} by the c -axis oscillation of in-plane O atoms in a single-layer model to explain the superconductive phonon anomalies in the important and well-characterised B -symmetric modes of the YBCO plane, as well as to obtain qualitative agreement with the results for other materials classes, with the observation of the spin gap in underdoped YBCO compounds, and with isotope effects at optimal doping. The reader is referred to this paper for the background to this section.

In Fig. 8(a) are shown the four c -axis modes of planar O atoms in the bilayer system: at the single-layer level [19], there is no difference between the 340cm^{-1} B_{1g} mode and the 193cm^{-1} B_{2u} mode, or between the 440cm^{-1} A_{1g} mode and the 307cm^{-1} A_{2u} mode. While this may be realistic for the B -symmetric modes, which develop no net dipole moment, it was found not to afford an acceptable description of either the A_{2u} mode, where there will be significant interplane charge motion during the phonon oscillation, or the A_{1g} mode, in which charge may be transferred out of the bilayer unit to neighbouring planes. Here we investigate the phonon anomalies in the bilayer in the presence of finite interplane coupling and superexchange, which as above lead to a finite interplane uniform spin order parameter, but no interplane singlet order. There have been indications within a phenomenological weak electron-phonon coupling model [48] that interlayer hopping terms are sufficient to reproduce the phonon anomalies measured [49] in the infrared-active A_{2u} mode.

Following Ref. [19], in the bilayer it is necessary to recast the phonon-spinon coupling vertices in terms of the diagonalising quasiparticle operators. Each operator f_k^α is a sum of 4 operators with coefficients given by (5), so that each vertex is a sum of 16 terms, each involving a contribution from both spin orientations. For the “uniform RVB” vertex one obtains on substitution

$$\begin{aligned} \sum_{\sigma} \bar{\chi} f_{k\sigma}^{1(2)\dagger} f_{k\sigma}^{1(2)} &= \frac{1}{2} \left[C_1 \gamma_{1k}^\dagger \gamma_{1k} + S_1 \gamma_{1k}^\dagger \gamma_{1-k}^\dagger \mp C_{12} \gamma_{1k}^\dagger \gamma_{2k} \mp S_{12} \gamma_{1k}^\dagger \gamma_{2-k}^\dagger \right. \\ &\quad + S_1 \gamma_{1-k} \gamma_{1k} + C_1 \gamma_{1-k} \gamma_{1-k}^\dagger \mp S_{12} \gamma_{1-k} \gamma_{2k} \pm C_{12} \gamma_{1-k} \gamma_{2-k}^\dagger \\ &\quad \mp C_{12} \gamma_{2k}^\dagger \gamma_{1k} \mp S_{12} \gamma_{2k}^\dagger \gamma_{1-k}^\dagger + C_2 \gamma_{2k}^\dagger \gamma_{2k} + S_2 \gamma_{2k}^\dagger \gamma_{2-k}^\dagger \\ &\quad \left. \mp S_{12} \gamma_{2-k} \gamma_{1k} \pm C_{12} \gamma_{2-k} \gamma_{1-k}^\dagger + S_2 \gamma_{2-k} \gamma_{2k} - C_2 \gamma_{2-k} \gamma_{2-k}^\dagger \right], \end{aligned} \quad (37)$$

and for the “singlet-RVB” vertex

$$\begin{aligned} (\Delta_k^* f_{k\uparrow}^{1(2)} f_{k\downarrow}^{1(2)} + \text{h.c.}) &= \frac{1}{2} \left[-S_1 \gamma_{1k} \gamma_{1k}^\dagger + C_1 \gamma_{1k} \gamma_{1-k} \pm S_{12} \gamma_{1k} \gamma_{2k}^\dagger \mp C_{12} \gamma_{1k} \gamma_{2-k} \right. \\ &\quad + C_1 \gamma_{1-k}^\dagger \gamma_{1k} + S_1 \gamma_{1-k}^\dagger \gamma_{1-k} \mp C_{12} \gamma_{1-k}^\dagger \gamma_{2k}^\dagger \mp S_{12} \gamma_{1-k}^\dagger \gamma_{2-k}^\dagger \\ &\quad \pm S_{12} \gamma_{2k} \gamma_{1k}^\dagger \mp C_{12} \gamma_{2k} \gamma_{1-k} - S_2 \gamma_{2k} \gamma_{2k}^\dagger + C_2 \gamma_{2k} \gamma_{2-k} \\ &\quad \left. \mp C_{12} \gamma_{2-k}^\dagger \gamma_{1k}^\dagger \mp S_{12} \gamma_{2-k}^\dagger \gamma_{1-k} + C_2 \gamma_{2-k}^\dagger \gamma_{2k} + S_2 \gamma_{2-k}^\dagger \gamma_{2-k} \right], \end{aligned} \quad (38)$$

where C_α and S_α denote $\cos 2\theta_k^\alpha$ and $\sin 2\theta_k^\alpha$, and C_{12} and S_{12} are the combinations $c_1 c_2 - s_1 s_2$ and $c_1 s_2 + s_1 c_2$, where c_α and s_α denote $\cos \theta_k^\alpha$ and $\sin \theta_k^\alpha$. The complexity contained in 2×2 matrix vertices and Nambu propagators for the $\vec{f}_k$ problem is transferred solely to the 4×4 matrix vertex in the quasiparticle problem, conserving the number of degrees of freedom.

In the even modes (A_{1g} and B_{1g}), the motions of the O atoms are out of phase between the planes, so that the effect of their motion on t and J is the same in each plane. Defining the phonon coordinate as positive when the oscillating O atom moves towards the plane of the Cu atoms (Fig. 8(b)), one has in obvious notation $u_x^1 = u_y^1 = u_x^2 = u_y^2 = u$ for A_{1g} and $u_x^1 = -u_y^1 = u_x^2 = -u_y^2 = u$ for B_{1g} . Thus in both cases the effective vertex from summing the contributions of both planes in (37) and (38) above is to cancel all of the terms mixing the indices 1 and 2 for both x and y bonds, leaving only those terms related to intraband processes. Conversely, the odd modes A_{2u} ($u_x^1 = u_y^1 = -u_x^2 = -u_y^2 = u$) and B_{2u} ($u_x^1 = -u_y^1 = -u_x^2 = u_y^2 = u$) will have vertices coupling only to interband scattering processes. This very simple but profound separation of the contributions may be understood readily, in that the even modes produce a pattern of modulations of t and J which is even in the unit cell, leading only to intraband transitions, and conversely for the pattern of modulations arising from the odd modes. The lowest-order quasiparticle polarisation terms contributing to the phonon self-energy may then be calculated to deduce the frequency shift and linewidth broadening of each type of mode. For the even modes (intraband), the contribution from normal scattering processes will vanish in the limit $q \rightarrow 0$, and one may follow [19] exactly to obtain

$$\delta\omega = c(\lambda_J J)^2 \frac{2}{N} \sum_k F_k^+ \frac{1}{\omega^2 - (2E_k^+)^2} \frac{\tanh\left(\frac{E_k^+}{2T}\right)}{E_k^+} + F_k^- \frac{1}{\omega^2 - (2E_k^-)^2} \frac{\tanh\left(\frac{E_k^-}{2T}\right)}{E_k^-}. \quad (39)$$

Here $c = \left(\frac{3}{4a}\right)^2 \langle u^2 \rangle = 1.18 \times 10^{-4}$, $\langle u^2 \rangle = \frac{\hbar}{2M\omega_0} = (0.055\text{\AA})^2$, in which $\omega_0 = 340\text{cm}^{-1}$ is taken from the B_{1g} phonon frequency and M is the mass of the O atom, and $E_k^\pm$ are given by (8). The form factors $F_k^\pm$ are

$$\begin{aligned} F_k^\pm &= 2\Delta^2 \left[\gamma_k (\xi_k \pm |M_k|) + \frac{3J}{4} \bar{\chi} \eta_k^2 \right]^2 & B_{1g} \\ F_k^\pm &= 2\Delta^2 \eta_k^2 \left[(\xi_k \pm |M_k|) + \frac{3J}{4} \bar{\chi} \gamma_k \right]^2 & A_{1g}, \end{aligned} \quad (40)$$

where $\gamma_k = \cos k_x + \cos k_y$, $\eta_k = \cos k_x - \cos k_y$, $\Delta = \langle \Delta_{ij} \rangle$ and $\bar{\chi} \equiv \langle \chi_{ij} \rangle + \frac{2t\delta}{3J}$. In this calculation we have not included an ‘‘interplane uniform’’ vertex with contributions proportional to B'_k and χ'_k , and will discuss the effects of such terms in detail below. The linewidth broadening can be computed from an analogous expression for the imaginary part of the self-energy.

For the odd modes, the interband scattering expression and form factors become

more complicated. One obtains

$$\begin{aligned} \delta\omega = & c(\lambda_J J)^2 \frac{1}{N} \sum_k F_k^n \frac{2(E_k^+ - E_k^-)}{\omega^2 - (E_k^+ - E_k^-)^2} (f(E_k^-) - f(E_k^+)) \\ & + F_k^a \frac{2(E_k^+ + E_k^-)}{\omega^2 - (E_k^+ + E_k^-)^2} (1 - f(E_k^-) - f(E_k^+)), \end{aligned} \quad (41)$$

in which the form factor for normal scattering events is

$$\begin{aligned} F_k^n &= \left[\sqrt{2} \Delta \gamma_k (c_1 s_2 + s_1 c_2) + \bar{\chi} \eta_k (c_1 c_2 - s_1 s_2) \right]^2 & B_{2u} \\ F_k^n &= \left[\sqrt{2} \Delta \eta_k (c_1 s_2 + s_1 c_2) + \bar{\chi} \gamma_k (c_1 c_2 - s_1 s_2) \right]^2 & A_{2u}, \end{aligned} \quad (42)$$

and that for anomalous scattering events is

$$\begin{aligned} F_k^a &= \left[\sqrt{2} \Delta \gamma_k (c_1 c_2 - s_1 s_2) + \bar{\chi} \eta_k (c_1 s_2 + s_1 c_2) \right]^2 & B_{2u} \\ F_k^a &= \left[\sqrt{2} \Delta \eta_k (c_1 c_2 - s_1 s_2) + \bar{\chi} \gamma_k (c_1 s_2 + s_1 c_2) \right]^2 & A_{2u}. \end{aligned} \quad (43)$$

It is straightforward to show that in the limit where the bands become degenerate, both sets of expressions reduce to the result expected from [19] for the single layer.

In Figs. 9-12 we show the phonon frequency shift $\delta\omega$ calculated using the above formulae, and the linewidth broadening $\delta\Gamma$, which is calculated from analogous expressions. We compute these anomalies only for the YBCO system, where the buckling of the O atoms in the bilayer structural unit is uniformly directed along the c -axis towards the symmetry plane of the unit cell, an arrangement which may yield constructive combination of the vertex contributions to give a linear spin-phonon coupling term in cases where the symmetries of the phonon distortion and the superconducting gap are compatible. By contrast, in BSCCO the single layer is orthorhombically buckled, into a tilting distortion which is odd in the unit cell [47], so there is no linear coupling term and the anomalies are expected to be negligibly small [19], as confirmed by experiment [50].

In Fig. 9(a) is shown $\delta\omega$ as a function of frequency for the g -symmetric modes, at $q = 0$ and at a temperature well below the singlet ordering transition T_{RVB} (Fig. 4). As in the single-layer case, low-frequency B -symmetric phonons will be softened by the spinon contribution, while high-frequency modes will show hardening. The characteristic frequency of the sign change is set by the value of the gap Δ_k near the heavily-contributing $(\pi, 0)$ points for the intraband processes in question. Quantitatively, it is somewhat smaller than in the single-layer solution, requiring that the analogue of the maximally-shifted B_{1g} mode in this system appear at $\omega \simeq 0.2J$, in comparison with the experimental mode frequency $\omega_0 \simeq 0.3J$. The imaginary part (Fig. 10(a)) has the expected form of a peak at the frequency where the real part changes sign, and shows that only phonon modes with frequencies close to this value will be significantly broadened.

For even, A -symmetric modes one finds also the result of the single-layer system, that the anomalies are negligible. The symmetry effects which cause the linear contribution to be small in this case are not affected by the introduction of interplane terms.

For odd (u -symmetric) modes one sees a different feature (Figs. 9(b), 10(b)). Because these couple only to interplane scattering processes, the characteristic frequency which determines the change from phonon softening to hardening is set not by twice the gap Δ_k , but by twice the interband energy separation $M_k \simeq \delta t_{\perp k}$, again at the $(\pi, 0)$ points where it is largest. From the results of Ref. [32] which were used in setting the interplane hopping parameter $t_{\perp}^0$, and are consistent with available experimental evidence, this separation is of order J , and so appears on an altogether different energy scale. Thus all u -symmetric phonons in the usual energy range will soften at the singlet ordering transition within such a model, and the linewidth broadening will be negligible.

In Figs. 11(a) and 12(a) are shown the temperature dependences of $\delta\omega$ and $\delta\Gamma$ for the g -symmetric modes; this is the quantity measured by experiment [51, 52], and should be directly comparable with the results for the B_{1g} and A_{1g} modes. One sees again that the results are very similar to the single-layer case: for the B_{1g} mode there is a gradual development of the frequency shift over a range of T below its onset at T_{RVB} , and a peak in the linewidth broadening there, both in good agreement with experiment [51]. The chosen mode frequency here is $\omega_0 = 0.2J$, which is somewhat below the experimental value, but is consistent with the location of the B_{1g} phonon being that for a maximal anomalous effect, and remains in a reasonable degree of correspondence with the physical system given the nature of the model. For the A_{1g} mode, the chosen frequency should be in the phonon hardening regime, but we show again $\omega_0 = 0.2J$ to emphasise that there is very little effect, and we comment on this in detail below. In Fig. 11(b) is the T dependence of $\delta\omega$ for a mode of frequency $\omega_0 = 0.2J$ and B_{2u} symmetry. Here ω_0 is so far from the characteristic frequency of the changeover from softening to hardening of u -symmetric phonons (Fig. 9(b)) that the development of the anomalous frequency shift is effectively immediate at T_{RVB} , as seen less obviously in the single layer model, and in good agreement with neutron scattering studies of this mode in $\text{YBa}_2\text{Cu}_3\text{O}_7$ [53]. Similarly, $\delta\Gamma$ for such phonons as a result of coupling to the spin is negligible within this model.

We have considered only $q = 0$ phonons here, for comparison with the results of Raman and infrared spectroscopy. For finite in-plane wavevector $\mathbf{q}$, one may expect the results to be similar to the single-layer case [19] for both g - and u - symmetric modes, namely that the anomalies show some features of the Fermi surface shape, and fall to zero as $\mathbf{q}$ approaches the $(\pi, 0)$ and (π, π) points. However, at finite q_z the intra- and interband scattering processes become mixed, so that both mode symmetries may show both characteristic energy scales. Of particular note is the situation at $\mathbf{q} = 0$, $q_z = \pi$, where the anomalies will remain large and the contributing processes will be completely reversed in (intra- or interband) nature. Thus it should be possible by neutron scattering to see that the onset of the frequency shift $\delta\omega$ for the B_{1g} phonon is very much sharper in temperature, if smaller in overall magnitude, as its frequency will no longer be close

to the relevant energy scale. Similarly, no alterations in the linewidth of this phonon should be detectable.

We have shown that the bilayer system retains a good description of the main qualitative features of the B -symmetric modes of planar O in the bilayer structural unit, although we stress again that the g - and u -symmetric modes have very different characteristic energy scales in the coupled bilayer system. However, the current level of treatment does not contain the physics required to explain the experimental results for the A -symmetric modes. Here we have included the interlayer hopping only through the way in which it arises in the quasiparticle transformation, and in this sense $t_{\perp k}$ is treated on an equal footing with the terms t'_k and t''_k , which are taken to determine $E_k^{\pm}$ but not to contribute qualitatively to the phonon anomalies [19]. This is also the approach taken by Ref. [48], but these authors chose the value of $t_{\perp}^0$ to coincide with the phonon frequency, in order to enhance the anomaly in their approach, and in addition used both less restrictive symmetry constraints and an isotropic gap function Δ_0 .

We have neglected the possibility of a contribution from modulation of the magnitude of the parameter $t_{\perp}^0$, because in none of the phonon modes considered do the Cu atoms of the bilayer change their relative separation. If one wishes to include such a modulation, for example on the grounds that additional transfer may occur through the neighbouring O atoms (whose separation changes in g -symmetric modes), it would appear as a term $\lambda_{\perp} \chi'_k (f_k^{1\dagger} f_k^2 + \text{h.c.})$ in the Hamiltonian. Substitution of the quasiparticle operators yields the same type of vertex terms as in the expressions (37) and (38), but with considerably smaller coefficient for reasonable values of the interplane phonon coupling $\lambda_{\perp}$. Thus the additional term would neither make a significant contribution nor appear with a symmetry factor which combines constructively with A -symmetric phonons. For this last reason, even a very strong modulation of the coefficient $t_{\perp}^0$ would not explain the experimental observations within the current theory. Such a modulation may in fact be present in the A_{2u} mode, where a very strong dipole moment is developed during the oscillatory motion, but it appears that this must be described by additional physics, related to the strong effects of the motion on interlayer charge transfer. Similarly, in the A_{1g} mode one must also take account of charge transfer, which may occur not between the planes of the bilayer but between each plane and the neighbouring BaO layers (in which O(4) is the apical oxygen atom of planar Cu(2)). Recent Raman-scattering experiments [54] suggest that the 440cm^{-1} A_{1g} planar mode and the 500cm^{-1} A_{1g} mode of apical O behave as split levels of a single, fundamental phonon frequency, indicating very strong coupling between in- and out-of-plane modes of these symmetries.

We conclude this section by noting that apart from the change in the nature of the u -symmetric modes discussed in detail above, none of the qualitative features extracted from the single-layer model [19] are altered in the bilayer system. In particular, the important result that the spin gap in the low-doping regime of the model is observable due to spin-phonon coupling [55] is retained, a point which reflects that the current model does not contain a difference in such properties between mono- and bilayer oxide materials. The isotope effect arising as a result of the coupling between the lattice and

the spin sector, which is responsible for superconductivity, is identical to that deduced from the single-layer model, because the bilayer treatment also does not reproduce the anomalous contributions from A -symmetric modes.

5 Summary

In summary, we have analysed the spin properties of a bilayer CuO_2 system, using the mean-field, slave-boson treatment of the extended t - J model. We find that the existence of d -symmetric spin singlet order in the plane, favoured by the single-layer model, acts to suppress interplane order completely below a characteristic, finite value of the interplane coupling. At this coupling, there is a very rapid crossover to a state of pure interplane singlet spin order, suggesting that the two types do not coexist. Working within the physical regime of interplane coupling, which mandates exclusively in-plane singlet pairing, we show that the observed Fermi surfaces of both BSCCO- and YBCO-like bilayer systems may be reproduced with realistic extended in-plane transfer integrals.

For each type of system, we have calculated two microscopic and physically observable properties related to the spin sector. The dynamic susceptibility $\chi(\mathbf{q}, \omega)$, computed at the RPA level, shows the spin fluctuation spectrum at all frequencies and wavevectors; the commensuration and excitation properties may be explained simply in terms of Fermi-surface scattering processes, but the remaining discrepancies between the theory and experiment in YBCO reveal that the problem is significantly more interesting and complex than mean-field physics. Anomalies in the frequency and linewidth of phonon modes in the YBCO system show a strong contrast between even (g -symmetric) modes, which couple to intraband scattering processes, and odd (u -symmetric) modes, which couple only to interband scattering. The model provides a reasonable account of the B -symmetric modes, which have no net electric dipole moment in each layer and so are not accompanied by charge motion, but cannot describe the A -symmetric modes, which do not satisfy this condition.

We conclude by commenting that because in this model the physical parameter regime remains dominated by the d -symmetric in-plane singlet order of the single-layer case, the results for observable quantities in the bilayer system differ qualitatively from those of the single layer in only a small number of features. We have carried out semi-quantitative calculations which may be compared in detail with experiment. This makes it possible to quantify the strengths and limitations of the current t - J model formulation, to learn which properties appear due to the bilayer nature, and to gain some evaluation of the restrictions of the mean-field theory and RPA approaches. Thus we may deduce the directions in which to look for the additional physics which will be required for a more complete description of the high- T_c systems. As an example of this process, we have assumed the existence of coherent single-particle transfer between the planes of the bilayer, and investigated the experimental consequences. The interlayer pair tunnelling theory [25] argues that such transfer cannot be coherent, and leads to some very different

physical predictions, which remain to be computed for similarly detailed comparison. Continuing rapid improvements in crystal quality, and in a variety of experimental techniques, will provide crucial evidence in favour of one of these viewpoints in the near future.

Acknowledgements

We are grateful to S. Maekawa for sharing results prior to publication. This work was supported financially by the Monbusho International Scientific Research Program No. 05044037 and the Grant-in-Aid for Scientific Research No. 04240103 of the Ministry of Education, Science and Culture, Japan. B.N. wishes to acknowledge the support of the Japan Society for the Promotion of Science.

References

- [1] A. J. Millis and H. Monien, Phys. Rev. Lett. **70**, 2810 (1993).
- [2] T. M. Rice, in The Physics and Chemistry of Oxide Superconductors, edited by Y. Iye and H. Yasuoka (Springer Verlag, Berlin, 1992); H. Fukuyama, Prog. Theor. Phys. Suppl. **108**, 287 (1992).
- [3] H. Yasuoka, T. Imai and T. Shimizu, in Strong Correlations and Superconductivity, edited by H. Fukuyama, S. Maekawa and A. P. Malozemoff (Springer Verlag, Berlin, 1989).
- [4] P. W. Anderson, Science **235**, 1196 (1987).
- [5] F.-C. Zhang and T. M. Rice, Phys. Rev. B **37**, 3759 (1988).
- [6] G. Baskaran, Z. Zou and P. W. Anderson, Solid State Commun. **63**, 973 (1987).
- [7] C. Gros, R. Joynt and T. M. Rice, Phys. Rev. B **36**, 8190 (1987).
- [8] G. Kotliar and J. Liu, Phys. Rev. B **38**, 5142 (1988).
- [9] Y. Suzumura, Y. Hasegawa and H. Fukuyama, J. Phys. Soc. Jpn. **57**, 2768 (1988).
- [10] T. Tanamoto, K. Kuboki and H. Fukuyama, J. Phys. Soc. Jpn. **60**, 3072 (1991).
- [11] T. Tanamoto, H. Kohno and H. Fukuyama, J. Phys. Soc. Jpn. **62**, 717 (1993).
- [12] Q. Si, Y. Zha, K. Levin, J. P. Lu and J. H. Kim, Phys. Rev. B **47**, 9055 (1993).
- [13] M. Lercher and J. Wheatley, Phys. Rev. B **49**, 736 (1994).
- [14] M. Lavagna and G. Stemann, Phys. Rev. B **49**, 4235 (1994).

- [15] K. Maki and H. Won, Phys. Rev. Lett. **72**, 1758 (1994).
- [16] M. Ubbens and P. A. Lee, Phys. Rev. B **50**, 438 (1994).
- [17] T. Tanamoto, H. Kohno and H. Fukuyama, J. Phys. Soc. Jpn. **62**, 1455 (1993) and **63**, 2741 (1994); H. Fukuyama, H. Kohno and T. Tanamoto, J. Low Temp. Phys. **95**, 309 (1994).
- [18] N. Nagaosa and P. A. Lee, Phys. Rev. Lett. **60**, 2450 (1990); P. A. Lee and N. Nagaosa, Phys. Rev. B **46**, 5621 (1992).
- [19] B. Normand, H. Kohno and H. Fukuyama, SISSA-Trieste Preprint No. 9502019.
- [20] L. B. Ioffe, A. I. Larkin, A. J. Millis and B. L. Altschuler, JETP Lett. **59**, 65 (1994).
- [21] A. I. Liechtenstein, I. I. Mazin and O. K. Andersen, Phys. Rev. Lett. **74**, 2303 (1995).
- [22] R. Eder, Y. Ohta and S. Maekawa, preprint.
- [23] A. J. Millis and H. Monien, Phys. Rev. B **50**, 16606 (1994).
- [24] A. J. Millis, L. B. Ioffe and H. Monien, J. Phys. Chem. Solids, to appear.
- [25] S. Chakravarty, A. Sudbø, P. W. Anderson and S. Strong, Science **261**, 337 (1993).
- [26] D. S. Dessau, Z.-X. Shen, D. M. King, D. S. Marshall, L. W. Lombardo, P. H. Dickinson, A. G. Loeser, J. DiCarlo, C.-H. Park, A. Kapitulnik and W. E. Spicer, Phys. Rev. Lett. **71**, 2781 (1993).
- [27] H. Ding, J. C. Campuzano, A. Bellman, T. Yokoya, M. R. Norman, M. Randeria, T. Takahashi, H. Katayama-Yoshida, T. Mochiku, K. Kadowaki and G. Jennings, preprint.
- [28] K. Gofron, J. C. Campuzano, A. A. Abrikosov, M. Lindroos, A. Bansil, H. Ding, D. Koelling and B. Dabrowski, Phys. Rev. Lett. **73**, 3302 (1994).
- [29] K. Gofron et al., to be published.
- [30] I. Affleck and J. B. Marston, Phys. Rev. B **37**, 3774 (1988).
- [31] K. Kuboki and P. A. Lee, preprint.
- [32] O. K. Andersen, O. Jepsen, A. I. Liechtenstein and I. I. Mazin, Phys. Rev. B **49**, 4145 (1994).
- [33] J. M. Tranquada, P. M. Gehring, G. Shirane, S. Shamoto and M. Sato, Phys. Rev. B **46**, 5561 (1993).

- [34] J. Rossat-Mignod, L. P. Regnault, P. Bourges, C. Vettier, P. Burlet and J. Y. Henry, *Physica B* **186-189**, 1 (1993).
- [35] R. Stern, M. Mali, J. Roos and D. Brinkman, unpublished.
- [36] M. Grüninger, J. Münzel, A. Gaymann, A. Zibold, H. P. Geserich and T. Kopp, preprint.
- [37] Y. Itoh, Ph.D. thesis, University of Tokyo (1994).
- [38] K. Kuboki, private communication.
- [39] H. Matsukawa and H. Fukuyama, *J. Phys. Soc. Jpn.* **58** 3687 (1989) and **59** 1723 (1990); J. H. Jefferson, H. Eskes and L. F. Feiner, *Phys. Rev. B* **45**, 7959 (1992).
- [40] H. A. Mook, M. Yethiraj, G. Aeppli, T. E. Mason and T. Armstrong, *Phys. Rev. Lett.* **70**, 3490 (1993).
- [41] H. A. Mook, P. Dai, G. Aeppli, T. E. Mason, N. E. Hecker, J. A. Harvey, T. Armstrong, K. Salama and D. Lee, *Physica B*, to appear.
- [42] H. A. Mook *et al.*, unpublished.
- [43] B. Keimer *et al.*, unpublished.
- [44] T. E. Mason, G. Aeppli, S. M. Hayden, A. P. Ramirez and H. A. Mook, *Phys. Rev. Lett.* **71**, 919 (1993).
- [45] Y. Endoh, private communication.
- [46] E. Demler and S. C. Zhang, preprint.
- [47] see R. M. Hazen, in *Physical Properties of High Temperature Superconductors II*, edited by D. Ginsberg (World Scientific, Singapore, 1990).
- [48] G. Hastreiter, F. Forsthofer and J. Keller, *Solid State Commun.* **88**, 769 (1993).
- [49] A. P. Litvinchuk, C. Thomsen and M. Cardona, *Solid State Commun.* **83**, 343 (1993).
- [50] S. Sugai, unpublished.
- [51] E. Altendorf, X. K. Chen, J. C. Irwin, R. Liang and W. N. Hardy, *Phys. Rev. B* **47**, 8140 (1993).
- [52] N. Pyka, W. Reichardt, L. Pintschovius, G. Engel, J. Rossat-Mignod and J. Y. Henry, *Phys. Rev. Lett.* **70**, 1457 (1993).
- [53] H. Harashina, K. Kodama, S. Shamoto, M. Sato, K. Kakurai and M. Nishi, preprint.

[54] M. Käll, private communication.

[55] M. Käll, A. P. Litvinchuk, P. Berastegui, L.-G. Johansson and L. Börjesson, *Physica C* **225**, 317 (1994).

Figure Captions

Fig. 1: Schematic representation of the bilayer system, showing transfer integrals for quasiparticle hopping and spin superexchange interactions.

Fig. 2: Variation of singlet spin ordering temperature T_{RVB} with interplane hopping parameter $t_{\perp}^0$, at fixed $J_{\perp} = 0.1J$. The in-plane transfer integrals are $t = 4J$, $t' = -\frac{1}{6}t$ and $t'' = \frac{1}{5}t$.

Fig. 3: Bonding ($\circ$) and antibonding ($\times$) Fermi surfaces for bilayer systems with doping $\delta = 0.2$, shown in the first quadrant. (a) BSCCO-like system: parameters are chosen, following Ref. [26], to give one open and one closed Fermi surface, strong nesting of the latter (antibonding) surface, and band degeneracy in the ΓM direction. (b) YBCO-like system: parameters are chosen, following Refs. [32, 28, 29], to give open Fermi surfaces for both bands, but with that for the bonding band close to “closing”, and to lift the degeneracy in the ΓM direction.

Fig. 4: Self-consistent solutions for the singlet-RVB order parameter $\Delta(T)$. (a) BSCCO-like system. (b) YBCO-like system.

Fig. 5: Dynamic susceptibility calculated as a function of $\mathbf{q}$ in the $2d$ Brillouin zone at fixed frequency $\omega = 0.3J$, low temperature and $q_z = 0$ (intraband processes only). In our notation, the X point corresponds to $(\pi, 0)$ and is symmetrical with Y, while M denotes the point (π, π) . (a) BSCCO-like system: note strong incommensuration arising from nesting of the antibonding band. (b) YBCO-like system: note commensurate nature of the peak intensity.

Fig. 6: Dynamic susceptibility calculated as a function of $\mathbf{q}$ in the $2d$ Brillouin zone at fixed frequency $\omega = 0.3J$, low temperature and $q_z = \pi$ (interband processes only). (a) BSCCO-like system: note that intensity of interband processes is considerably higher than in the intraband case, and that they exhibit only minor incommensuration. (b) YBCO-like system: susceptibility remains commensurate at all q_z .

Fig. 7: Dynamic susceptibility calculated as a function of ω at fixed in-plane wavevector $\mathbf{q} = (\pi, \pi)$, with $q_z = 0$ ($\times$), $q_z = \pi/2$ ($\square$) and $q_z = \pi$ ($\circ$). (a) BSCCO-like system. The main peak is due to states pushed out of the gap, and is absent for the $q_z = 0$ case, primarily because the antibonding Fermi surface is everywhere smaller than (π, π) . The shoulder at $\omega \sim J$ indicates the energy separation above which there are no more states separated by (π, π) in $\mathbf{q}$ space for the BSCCO-like dispersion. (b) YBCO-like system. The two contributions to each peak are due to gaps with different values on each band at the wavevector $\mathbf{k}$ corresponding to (π, π) scattering. For the YBCO dispersion, the “shoulder” occurs at $\omega = 1.2J$.

Fig. 8: (a) Atomic motions and experimentally observed frequencies of the four phonon modes involving c -axis motion of in-plane O atoms in the buckled CuO_2 bilayer. (b) Phonon coordinate labelling convention.

Fig. 9: Phonon frequency shift $\delta\omega$ as a function of frequency and at $T = 0.05T_{RVB}$ in the YBCO-like system. (a) B_{1g} - ($\circ$) and A_{1g} -symmetric ($\times$) modes: the frequency of the sign-change is characterised by the gap value at the strongly-contributing $(\pi, 0)$ points, because only intraband processes are involved. (b) B_{2u} - ($\circ$) and A_{2u} -symmetric ($\times$) modes: because the phonon coordinate couples only to interband processes, the frequency of the sign-change is characterised by the value of the interplane hopping term at $(\pi, 0)$ points.

Fig. 10: Phonon linewidth broadening $\delta\Gamma$ at $q = 0$, as a function of frequency and at $T = 0.05T_{RVB}$ in the YBCO-like system. (a) B_{1g} - ($\circ$) and A_{1g} -symmetric ($\times$) modes. (b) B_{2u} - ($\circ$) and A_{2u} -symmetric ($\times$) modes.

Fig. 11: (a) Phonon frequency shift $\delta\omega$ as a function of T for hypothetical B_{1g} - ($\circ$) and A_{1g} -symmetric ($\times$) modes of frequency $\omega_0 = 0.2J$, at $q = 0$ for the YBCO-like system. Because the mode frequency is close to the characteristic frequency of the sign-change (Fig. 9(a)), the full frequency shift develops over a range of T . (b) $\delta\omega$ for B_{2u} - ($\circ$) and A_{2u} -symmetric ($\times$) modes of the same frequency. Because the ω_0 is far from the characteristic frequency of the sign-change (Fig. 9(b)), the onset is very abrupt.

Fig. 12: (a) Phonon linewidth broadening $\delta\Gamma$ as a function of T for hypothetical B_{1g} - ($\circ$) and A_{1g} -symmetric ($\times$) modes of frequency $\omega_0 = 0.2J$, at $q = 0$ for the YBCO-like system. Because ω_0 is close to the frequency of the peak (Fig. 10(a)), a significant broadening is observed. (b) $\delta\Gamma$ for B_{2u} - ($\circ$) and A_{2u} -symmetric ($\times$) modes the same frequency. Because ω_0 is so far from the characteristic frequency of the sign-change (Fig. 10(b)), the imaginary part is negligible.